



ELSEVIER

Thermochimica Acta 71135 (2002) 1–12

thermochimica
acta

www.elsevier.com/locate/tca

The enthalpies of formation of two dibenzocyclooctadienones

St. Perisanu^a, Iulia Contineanu^b, M.D. Banciu^c, Joel F. Liebman^d, Behzad S. Farivar^d,
Melissa A. Mullan^d, James S. Chickos^{e,*}, Nigam Rath^e, Dorothea M. Hillesheim^e

^a Laboratory of General Chemistry, Polytechnic University of Bucharest, Str. Polizu nr. 1, Bucuresti 78126, Romania

^b Institute of Physical Chemistry, Romanian Academy, Spl. Independentei nr. 202, Bucuresti 77208, Romania

^c Laboratory of Organic Chemistry, Polytechnic University of Bucharest, Spl. Independentei nr. 313, Bucuresti 76206, Romania

^d Department of Chemistry and Biochemistry, University of Maryland, Baltimore County, Baltimore, MD 21250, USA

^e Department of Chemistry, University of Missouri-St. Louis, St. Louis, MO 63121, USA

Received in revised form 16 September 2002; accepted 20 September 2002

Abstract

The standard molar enthalpies of formation ($\Delta_f H_m^0$ (s)/kJ mol⁻¹) for 2,3:6,7-dibenzocycloocta-2,6-dien-1-one and 2,3:7,8-dibenzocycloocta-2,7-dien-1-one [6*H*-11,12-dihydro-dibenzo[*a,e*]cycloocten-5-one (ketone **1**) and 10*H*-11,12-dihydrodibenzo[*a,d*]-cycloocten-5-one (ketone **2**), respectively] were derived from enthalpies of combustion, measured by means of a microbomb calorimeter. The fusion and vaporization enthalpies of these compounds were obtained from DSC and correlation gas chromatography measurements. The standard molar enthalpies of formation in the gas phase were calculated by combining the condensed phase standard molar enthalpies of formation with the fusion and vaporization enthalpies adjusted to 298.15 K. Values for $\Delta_f H_m^0$ (g) of (-39.9 ± 5.5) and (-14.8 ± 5.3) kJ mol⁻¹ were obtained for 2,3:6,7-dibenzocycloocta-2,6-dien-1-one and 2,3:7,8-dibenzocycloocta-2,7-dien-1-one, respectively. Quantum chemical calculations are reported for the compounds investigated experimentally and an additional four isomers. Isomerization enthalpies are derived from computed energies. The enthalpies of formation are also calculated by group additivity, compared with the experimental values and then correlated with the structure of the molecules investigated. The X-ray analysis of ketone **1** is also reported.


© 2002 Published by Elsevier Science B.V.

Keywords: Dibenzocyclooctadienones; Molar enthalpies; DSC

1. Introduction

This paper continues the thermochemical investigation of some dibenzoannelated cycloalkane derivatives. In previous papers, the standard enthalpies of formation of some dibenzocycloheptane ketones [1] and of some dibenzocycloheptane alcohols [2] were measured.

With the aim of a comparative study of the enthalpies of formation of dibenzocycloalkane derivatives, data on the standard molar enthalpies of formation of the two dibenzocyclooctadienones, ketones **1** and **2** are added in this paper:




2,3:6,7-dibenzocycloocta-2,6-dien-1-one (ketone **1**)


* Corresponding author. Tel.: +1-314-516-5377;

fax: +1-314-516-5342.


E-mail address: jsc@umsl.edu (J.S. Chickos).



2,3:7,8-dibenzocycloocta-2,7-dien-1-one (ketone 2)




2,3:4,5-Dibenzocycloocta-2,4-dien-1-one (ketone 6)




2,3:5,6-dibenzocycloocta-2,5-dien-1-one (ketone 3)

40 During the course of our investigations of the thermal properties of ketones **1** and **2**, some concern was
 41 expressed regarding the structure of ketone **1**. A question arose regarding whether the structure reported as
 42 ketone **1** was actually ketone **3**. The ^1H and ^{13}C NMR spectra of **1** were not particularly revealing. The formation of ketone **1** requires two ring expansions and
 43 the possibility of other rearrangements competing with ring expansion could not be conclusively eliminated.
 44 As a result, the structure of ketone **1** was unambiguously determined by X-ray crystallography. This paper reports the structure of ketone **1** and thermochemical
 45 properties of ketones **1** and **2**. These results are compared to ab initio quantum mechanical calculations for ketones **1–3** as well as for three additional isomers,
 46 ketones **4–6**.



3,4:6,7-dibenzocycloocta-3,6-dien-1-one (ketone 4)



3,4:5,6-Dibenzocycloocta-3,5-dien-1-one (ketone 5)

55

2. Experimental

56

2.1. Synthesis and purification

57

Ketone **1** was obtained using procedures described in previous papers [3,4]. The buffered acetolysis of ditosylate **7** afforded the alcohol **8** in high yield. Catalytic hydrogenation of **8** gave rise to the saturated alcohol **9**, which was subsequently oxidized to ketone **1** (Scheme 1).

58

59

60

61

62

63

The synthesis of ketone **2** was performed using a reaction sequence described in the literature [5] (Scheme 2).

64

65

66

The spectral data (IR, ^1H NMR, ^{13}C NMR and MS) of ketones **1** (mp = 368 K, after two recrystallizations from ethanol) and **2** (mp = 421 °C, after two recrystallizations from 2-propanol) have previously been reported [5,6]. Samples analyzed by gas chromatography to be over 99.9% pure were used in the subsequent studies.

67

68

69

70

71

72

73

2.2. X-ray crystal structure

74

Colorless crystals of ketone **1** were grown by slow evaporation from ethanol solution at ambient temperature. A crystal with dimensions 0.34 mm × 0.16 mm × 0.10 mm was mounted on a glass fiber in a random orientation. Preliminary examination and data collection was performed using a Bruker SMART Charge Coupled Device (CCD) Detector system single crystal X-ray diffractometer using graphite monochromated Mo K α radiation ($\lambda = 0.71073 \text{ \AA}$) equipped with a sealed tube X-ray source at 170 K. Preliminary unit cell constants were determined with a set of 45 narrow frames (0.4° in ϖ) scans. The data set collected consisted of 3636 frames with a frame width of 0.3° in ϖ and counting time of 15 s/frame at a crystal to detector distance of 4.900 cm. The double pass method of

75

76

77

78

79

80

81

82

83

84


85

86

87

88

89




Scheme 1.

90 scanning was used to exclude any noise. The collected
 91 frames were integrated using an orientation matrix de-
 92 termined from the narrow frame scans. SMART and
 93 SAINT software packages (Bruker Analytical X-ray,
 94 Madison, WI, 2001) were used for data collection and
 95 data integration. Analysis of the integrated data did
 96 not show any decay. Final cell constants were deter-
 97 mined by a global refinement of xyz centroids of 4048
 98 reflections ($\theta < 28.3^\circ$). Collected data were corrected
 99 for systematic errors using SADABS [7] based on the
 100 Laue symmetry using equivalent reflections. The inte-
 101 gration process yielded 11,712 reflections of which
 102 1592 were independent reflections.

103 Crystal data and intensity data collection param-
 104 eters are listed in Table 1. Structure solution and refine-
 105 ment were carried out using the SHELXTL software
 106 package (Sheldrick, G.M., Bruker Analytical X-ray
 107 Division, Madison, WI, 2001). The structure was
 108 solved by Direct methods and refined successfully in

the space group $C2/c$. Full matrix least-squares refine- 109
 ment was carried out by minimizing $\sum w(F_O^2 - F_C^2)^2$. 110
 The non-hydrogen atoms were refined anisotropically 111
 to convergence. The hydrogen atoms were treated 112
 using appropriate riding model (AFIX m3). The final 113
 residual values were $R(F) = 4.5\%$ for 1176 obser- 114
 ved reflections [$I > 2\sigma(I)$] and $wR(F^2) = 11.6\%$; 115
 $s = 1.0$ for all data. The carbonyl O is disordered 116
 due to forced crystallographic symmetry. The O atom 117
 was refined with a 50% occupancy factor. Structure 118
 refinement parameters are listed in Table 1. A projec- 119
 tion view of ketone **1** with non-hydrogen atoms rep- 120
 resented by 50% probability ellipsoids, and showing 121
 the atom labeling is presented in Fig. 1. 122

Complete listings of the atomic coordinates for 123
 the non-hydrogen atoms and the geometrical param- 124
 eters, positional and isotropic displacement coeffi- 125
 cients for hydrogen atoms, anisotropic displacement 126
 coefficients for the non-hydrogen atoms are deposited 127



Scheme 2.

Table 1
Crystal data and structure refinements for ketone 1

Identification code	CCDC#181758
Empirical formula	C ₁₆ H ₁₄ O
Formula weight	222.27
Temperature (K)	170 (2)
Wavelength (Å)	0.71073
Crystal system	Monoclinic
Space group	C2/c
Unit cell dimensions (Å)	$a = 18.2308$, $\alpha = 90^\circ$ $b = 5.10800$ (10), $\beta = 117.934$ (1) $^\circ$ $c = 14.0319$ (3), $\gamma = 90^\circ$
Volume (Å ³ , z)	1154.45 (4), 4
Density (Mg/m ³)	1.279
Absorption coefficient (mm ⁻¹)	0.078
$F(000)$	472
Crystal size (mm)	0.34 × 0.16 × 0.10
θ range for data collection	2.53–28.27 $^\circ$
Limiting indices	$-24 \leq h \leq 21$, $-6 \leq k \leq 6$, $-18 \leq l \leq 18$
Reflections collected	11712
Independent reflections	1436 ($R_{\text{int}} = 0.038$)
Completeness to $\theta = 28.27^\circ$	100.0%
Absorption correction	None
Maximum and minimum transmission	0.9922 and 0.9740
Refinement method	Full matrix least squares on F^2
Data/restraints/parameters	1436/0/79
Goodness of fit on F	1.051
Final R indices ($I > 2\sigma(I)$)	$R1 = 0.0449$, $wR2 = 0.1083$
R indices (all data)	$R1 = 0.0567$, $wR2 = 0.1156$
Largest diff. peak and hole (eÅ ⁻³)	0.177 and -0.177




Fig. 1. A projection view of ketone 1 illustrating the forced crystallographic symmetry.

128 with the Cambridge Crystallographic Data Center
129 (CCDC#181758).

130 2.3. Fusion enthalpies

131 Fusion enthalpies were measured on a Perkin Elmer
132 DSC 7 using sealed stainless steel cells. The instru-
133 ment was calibrated using indium metal. Temperature
134 was scanned at 2 K/min. The results are reported in
135 Table 2. No other transitions were observed between
136 room temperature and the melting temperature.

137 2.4. Vaporization enthalpies

138 The vaporization enthalpies were derived by cor-
139 relation gas chromatography (CGC) using a Hewlett

140 Packard GC 5890 Series II. The column used for all
141 experiments was a HP1 (cross linked methyl silicone
142 gum) with the dimensions 12 m × 0.2 mm × 0.32 μm
143 (film thickness). A total of four mixtures have been
144 measured. All mixtures were prepared by dissolving
145 the respective compounds in methylene chloride. The
146 first mixture contained the reference compounds: ben-
147 zophenone, 9-heptadecanone and heptadecanoic acid

Table 2
Experimental fusion enthalpies by DSC

Compound	m (g)	T_{fus} (K)	$\Delta_{\text{fus}}H_m(T_{\text{fus}})$ (kJ mol ⁻¹)
Ketone 1			
Run 1	0.00236	366.3	17.40
Run 2	0.00161	364.1	17.62
Run 3	0.00192	366.2	16.84
Run 4	0.00199	368.6	16.70
Run 5	0.00218	367.9	17.52
Mean value		366.6	17.2 ± 0.5
Ketone 2			
Experiment 1	0.00486	420.1	28.03
Experiment 2	0.00506	420.1	27.12
Experiment 3	0.00413	419.7	28.28
Mean value		420.0	27.8 ± 0.7

Table 3
Some thermochemical properties of the vaporization standards

	$\Delta_{\text{vap}}H_{\text{m}}(T_{\text{m}}(\text{K}))$	$T_{\text{m}}(\text{K})$	$C_{\text{Pm}}(\text{l})(298.15\text{ K})(\text{J mol}^{-1}\text{ K}^{-1})$	$\Delta_{\text{vap}}H_{\text{m}}(298.15\text{ K})(\text{kJ mol}^{-1})$ [8]
Benzophenone	65.1	448	300.1	78.4
9-Heptadecanone	78.3	454	568.0	103.0
Methyl tetradecanoate	75.6	404	515.8	90.9
Methyl heptadecanoate	84.4	436	611.5	107.8
Methyl octadecanoate	83.2	442	643.4	108.8

148 methyl ester and the two ketones. The second mixture
149 contained the same reference compounds but only ke-
150 tone **1**. The third mixture contained the two ketones
151 and two more reference compounds, methyl tetradeca-
152 canoate and methyl octadecanoate. Ketone **1** exhib-
153 ited lower retention times than ketone **2**. The fourth
154 mixture was run as a control to demonstrate that the
155 vaporization enthalpies of the classes of compounds
156 used as standards correlate with each other. Vaporiza-
157 tion enthalpy data and estimated molar heat capacities
158 of the standards are detailed in Table 3 [8]. Eq. (1)
159 was used to adjust vaporization enthalpies from the
160 literature when necessary to 298.15 K:

$$\begin{aligned}
 162 \quad \Delta_{\text{vap}}H_{\text{m}}(298.15\text{ K}/\text{J mol}^{-1}) \\
 163 \quad &= \Delta_{\text{vap}}H_{\text{m}}(T_{\text{m}}) \\
 164 \quad &+ [10.58 + 0.26(C_{\text{Pm}}(\text{l})(298.15\text{ K})) \\
 165 \quad &\times [T_{\text{m}} - 298.15] \quad (1)
 \end{aligned}$$

166 The terms T_{m} and $C_{\text{Pm}}(\text{l})(298.15\text{ K})$ represent the
167 mean temperature of measurement and the heat ca-
168 pacity of the compounds at 298.15 K. The latter was
169 estimated by group additivity [9,10].

170 The experimental retention time data of methylene
171 chloride and the different compounds in each mixture
172 and the temperatures of each experiment are shown
173 in Table 4. The corrected retention times were calcu-
174 lated by subtracting the retention time of the methy-
175 lene chloride from the retention times of each com-
176 pound at each temperature. Methylene chloride is not
177 retained by the column under these conditions. Note
178 that the retention time of CH_2Cl_2 increases with in-
179 creasing temperature, contrary to the retention times
180 of the other components. The increase in the retention
181 time of CH_2Cl_2 with temperature is a consequence of
182 the corresponding increase in the viscosity of the He
183 carrier gas.

2.5. Enthalpies of combustion

184

185 The substances investigated were available in
186 small amounts and required the use of microcom-
187 bustion techniques. The enthalpies of combustion
188 were obtained using a two cell Tian–Calvet type
189 microcalorimeter that has been previously described
190 [11]. The calorimeter has two measuring elements
191 disposed symmetrically in a massive aluminium
192 block. The block is fastened inside of a thermostated
193 bath. The temperature of the water is maintained at
194 $298.15 \pm 0.01\text{ K}$. All around each calorimetric unit
195 there are 60 copel-nicrothal thermocouples. The ther-
196 mocouples were connected through a special switch,
197 designed to prevent spurious thermoelectrical forces,
198 to a Hewlett-Packard multimeter, Model 34401A. The
199 multimeter in turn was connected to a computer for
200 acquisition of the experimental data. The maximum
201 deviation (in mV) of the ballistic curves, which is
202 proportional to the areas that these delimitate, was
203 obtained by means of the Benchlynk program.

204 The microbombs constructed in the Laboratory of
205 Thermodynamics of the Institute of Physical Chem-
206 istry of Bucharest have also been described [12]. They
207 have been designed to burn samples that weigh less
208 than 8 mg. The microbombs were manufactured as
209 similarly as possible, to eliminate distortions originat-
210 ing from thermal transfer. The samples were pressed
211 into small discs 2.5 mm in diameter and about 1 mm
212 in height so that the mass of the sample did not exceed
213 8 mg. Ignition was carried out using cotton thread,
214 10 mm long, knotted on platinum wire and stretched
215 between the two electrodes. The heat of combustion of
216 the cotton thread was measured as $-16399 \pm 23\text{ J/g}$.
217 The contribution of the cotton thread is listed as q_c ,
218 the heat due to ignition in Table 5. Both the sample
219 and reference cell were flushed and filled with oxygen
220 to a final pressure of 3040 kPa. Both cells were ins-

Table 4
Gas chromatographic retention times t_r (min)

	T (K)						
	414.65	419.55	424.55	429.55	434.45	439.25	444.15
Mixture 1							
CH ₂ Cl ₂	0.215	0.220	0.219	0.219	0.222	0.222	0.226
Benzophenone	2.895	2.447	2.079	1.791	1.554	1.355	1.195
9-Heptadecanone	10.607	8.290	6.643	5.385	4.405	3.630	3.020
Methyl heptadecanoate	22.192	17.197	13.420	10.600	8.448	6.795	5.510
Ketone 1	10.607	8.610	7.010	5.785	4.815	4.040	3.417
Ketone 2	14.490	11.694	9.482	7.775	6.425	5.350	4.487
	414.65	419.55	424.55	429.55	434.45	439.35	444.05
Mixture 2							
CH ₂ Cl ₂	0.219	0.219	0.221	0.224	0.227	0.225	0.232
Benzophenone	2.912	2.457	2.091	1.799	1.558	1.362	1.197
9-Heptadecanone	10.515	8.318	6.673	5.412	4.420	3.648	3.030
Methyl heptadecanoate	22.225	17.231	13.471	10.645	8.469	6.819	5.521
Ketone 1	10.515	8.553	7.010	5.797	4.822	4.050	3.415
	414.9	419.7	424.6	429.6	434.5	439.1	444.3
Mixture 3							
CH ₂ Cl ₂	0.222	0.220	0.224	0.225	0.221	0.225	0.229
Benzophenone	2.915	2.457	2.093	1.799	1.553	1.355	1.198
Methyl tetradecanoate	5.279	4.266	3.493	2.893	2.409	2.028	1.731
9-Heptadecanone	10.556	8.323	6.681	5.415	4.413	3.630	3.034
Methyl heptadecanoate	22.283	17.262	13.502	10.659	8.481	6.795	5.531
Ketone 1	10.556	8.578	7.030	5.820	4.835	4.035	3.424
Ketone 2	14.510	11.700	9.525	7.819	6.448	5.355	4.503
Methyl octadecanoate	36.060	27.576	21.294	16.590	13.021	10.290	8.266
	414.9	419.7	424.6	429.6	434.4	439.2	444.1
Mixture 4							
CH ₂ Cl ₂	0.215	0.219	0.223	0.225	0.221	0.223	0.223
Benzophenone	2.902	2.452	2.091	1.796	1.554	1.352	1.188
2-Pentadecanone	4.54	3.717	3.07	2.56	2.15	1.818	1.555
Methyl tetradecanoate	5.238	4.256	3.489	2.887	2.408	2.022	1.718
2-Heptadecanone	11.832	9.372	7.493	6.036	4.911	4.014	3.319
Methyl hexadecanoate	13.683	10.771	8.548	6.844	5.53	4.49	3.689
Methyl heptadecanoate	22.16	17.195	13.449	10.611	8.449	6.766	5.479

221 erted into the calorimeter, initially at 298.15 ± 0.01 K
 222 and connected to the ignition device. Only the sample
 223 microbomb contained the cotton thread.

224 Ketone 2 did not compress well into pellets; con-
 225 sequently, paraffin film was used to seal the crucible.
 226 The enthalpy of combustion of the paraffin film was
 227 measured separately and was found to be $46,230 \pm$
 228 19 kJ g^{-1} . Ignition was triggered after introduction of
 229 the microbombs in the calorimetric block and a stable
 230 base line was established. A ballistic type curve was

recorded following ignition,

$$Q = kV \quad (2)$$

231 where k stands for the calibration constant of the
 232 calorimeter (J mV^{-1}) and V for the maximum devia-
 233 tion (mV). The calibration constant of the calorimeter
 234 was determined by means of benzoic acid combustion
 235 (standard reference material from NIST) and was cal-
 236 culated from the results of eight experiments. It was
 237 found to amount to $879.46 \pm 0.90 \text{ J mV}^{-1}$.
 238
 239

Table 5
Combustion data of ketone **1**^a

<i>m</i> (g)	<i>V</i> (mV)	<i>q_c</i> (J)	<i>m_{par}</i> (g)	<i>q_{par}</i> (J)	<i>Q</i> (J)	$\Delta\mu_{\text{exp}}$ (J/g)	$\Delta\mu_{\text{cor}}$ (J/g)
0.00391	0.258	4.88	0.00170	79.13	227.27	36641	36637
0.00475	0.660	6.18	0.00474	219.8	580.12	36628	36625
0.00410	0.336	6.83	0.00299	138.82	195.80	36623	36620
0.00469	0.673	3.25	0.00899	416.45	591.57	36646	36642
0.00445	0.303	4.72	0.00215	99.96	266.63	36642	36638
0.00373	0.262	7.81	0.0018	85.74	229.91	36632	36628
0.00366	0.220	4.88	0.00118	54.81	193.81	36644	36640
0.00326	0.249	6.83	0.00380	92.90	219.34	36695	36691
0.00620	0.419	7.48	0.00288	133.74	368.63	36683	36679
0.00657	0.398	5.52	0.00222	103.19	349.91	36711	36707

^a Average value $\Delta\mu_{\text{cor}} = -(36.651 \pm 0.019) \text{ kJ g}^{-1}$; $(8146.4 \pm 4.3) \text{ kJ mol}^{-1}$ (uncertainty ± 2 standard deviations of the mean); *m*: sample mass; *q_c*: heat due to ignition; *m_{par}*: mass of the paraffin film; *q_{par}*: heat of burning of the paraffin film; *Q*: total heat; $\Delta\mu_{\text{exp}}$: experimental specific energy of combustion; $\Delta\mu_{\text{cor}}$: corrected (Washburn) specific energy of combustion.

Table 6
Combustion data of ketone **2**^a

<i>m</i> (g)	<i>V</i> (mV)	<i>Q</i> (J)	<i>q_c</i> (J)	$\Delta\mu_{\text{exp}}$ (J g ⁻¹)	$\Delta\mu_{\text{corr}}$ (J g ⁻¹)
0.00622	0.260	228.38	4.55	36730	36726
0.00360	0.154	134.92	2.76	36703	36699
0.00704	0.310	265.33	6.98	36722	36718
0.00498	0.214	188.30	5.68	36699	36695
0.00641	0.276	242.44	6.98	36764	36760
0.00572	0.245	215.01	5.36	36739	36736
0.00603	0.257	225.92	4.62	36716	36712
0.00650	0.277	243.49	5.05	36694	36690

^a Average value $(-36.721 \pm 0.017) \text{ kJ g}^{-1}$; $(8161.9 \pm 3.7) \text{ kJ mol}^{-1}$; For a key, see footnote in Table 5.

240 The results of the combustion experiments are
241 shown in Tables 5 and 6. At least 10 runs were made
242 for each compound, but some of them were rejected
243 because of incomplete combustion. The data in the
244 tables come from experiments that gave reproducible
245 results with no evidence of soot present following
246 combustion. The reliability of the calorimeter was
247 confirmed by means of combustion of other reference
248 compounds such as salicylic acid and urea [12].
249 All uncertainties reported below are two standard
250 deviations unless otherwise noted.

251 3. Results

252 Condensed phase standard molar enthalpies of
253 combustion and formation of the two ketones are

summarized in Table 7. Values of -285.83 and $-393.51 \text{ kJ mol}^{-1}$ were used for the enthalpy of formation of $\text{H}_2\text{O}(\text{l})$ and $\text{CO}_2(\text{g})$ [13]. The standard molar enthalpies of sublimation of these two materials were obtained by combining the vaporization and fusion enthalpies adjusted to 298.15 K [14].

3.1. Vaporization enthalpies

Enthalpies of transfer from solution to the vapor, $\Delta_{\text{sln}}^{\text{g}} H_{\text{m}}(T_{\text{m}})$, were measured by correlation gas chromatography. They were calculated as the product of the gas constant, *R*, and the slope of the line obtained by plotting of the natural logarithm of reciprocal corrected retention time versus $1/T$ (K^{-1}) as previously reported [14]. The temperature *T_m* refers to the mean temperature of measurement. Correlating $\Delta_{\text{sln}}^{\text{g}} H_{\text{m}}(T_{\text{m}})$ with the vaporization enthalpies of the standards at 298.15 K, results in a linear relationship that can be used to obtain $\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K})$ of the unknowns. The enthalpies of transfer measured for the standards and for ketones **1** and **2** are summarized in Table 8. The correlation equations used to

Table 7
Standard enthalpies of combustion and formation of crystalline ketones **1** and **2** at 298.15 K

Compound	$-\Delta_{\text{c}} H^{\circ}$ (kJ mol ⁻¹)	$-\Delta_{\text{f}} H^{\circ}(\text{s})$ (kJ mol ⁻¹)
Ketone 1	8153.8 ± 4.3	143.2 ± 4.3
Ketone 2	8169.4 ± 3.7	127.6 ± 3.7

Table 8

A summary of the vaporization enthalpies obtained by correlation gas chromatography for mixtures 1–4 and the corresponding correlation equations

Compound	$\Delta_{\text{sln}}^{\text{g}} H_{\text{m}}(T_{\text{m}})$ (kJ mol ⁻¹)	$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K})$ literature (kJ mol ⁻¹)	$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K})$ calculated (kJ mol ⁻¹)
Mixture 1			
Benzophenone	52.66	78.4	79.1
9-Heptadecanone	67.75	103.0	100.7
Methyl heptadecanoate	73.86	107.8	109.4
Ketone 1	61.23		91.3
Ketone 2	62.68		93.4
$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K}) = (1.43 \pm 0.19) \Delta_{\text{sln}}^{\text{g}} H_{\text{m}}(T_{\text{m}}) + (3.64 \pm 2.9); \quad r^2 = 0.983$			
Mixture 2			
Benzophenone	53.09	78.39	79.3
9-Heptadecanone	67.43	102.99	100.2
Methyl heptadecanoate	73.93	107.8	109.7
Ketone 1	60.83		90.6
$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K}) = (1.46 \pm 0.23) \Delta_{\text{sln}}^{\text{g}} H_{\text{m}}(T_{\text{m}}) + (1.74 \pm 3); \quad r^2 = 0.975$			
Mixture 3			
Benzophenone	53.40	78.4	78.1
9-Heptadecanone	68.04	95.5	97.0
Methyl heptadecanoate	74.53	107.8	105.3
Methyl tetradecanoate	63.36	90.9	90.9
Methyl octadecanoate	78.16	108.8	110.0
Ketone 1	61.35		88.4
Ketone 2	63.05		90.5
$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K}) = (1.29 \pm 0.092) \Delta_{\text{sln}}^{\text{g}} H_{\text{m}}(T_{\text{m}}) + (9.43 \pm 1.8); \quad r^2 = 0.985$			
Mixture 4			
Benzophenone	53.62	78.4	76.2
2-Pentadecanone	61.70	83.7	87.4
Methyl tetradecanoate	63.50	90.9	89.9
2-Heptadecanone	69.28	95.8	98
Methyl hexadecanoate	71.12	101.3	100.5
Methyl heptadecanoate	74.90	107.8	105.8
$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K}) = (1.39 \pm 0.16) \Delta_{\text{sln}}^{\text{g}} H_{\text{m}}(T_{\text{m}}) + (1.55 \pm 2.69); \quad r^2 = 0.952$			

275 calculate the vaporization enthalpies are included in
 276 the table for each mixture. The equation of each line
 277 was generated by a linear regression analysis. Fig. 2
 278 illustrates the correlation observed for mixture 3. As
 279 noted above, the results from mixture 4 suggest that
 280 the enthalpies of transfer of the compounds chosen as
 281 standards correlate with each other.

282 3.2. Fusion enthalpy

283 The fusion enthalpy of ketones **1** and **2**, $\Delta_{\text{fus}} H_{\text{m}}(T_{\text{fus}})$,
 284 was adjusted from T_{fus} to 298.15 K using the follow-

ing equation [14,15].

$$\begin{aligned} \Delta_{\text{fus}} H_{\text{m}}(298.15 \text{ K}/\text{J mol}^{-1}) & \quad 287 \\ = \Delta_{\text{fus}} H_{\text{m}}(T_{\text{fus}}) + [0.75 + 0.15(C_{\text{Pm}}(\text{s})(298.15 \text{ K})) & \quad 288 \\ \times [T_{\text{fus}} - 298.15] - [10.58 & \quad 289 \\ + 0.26(C_{\text{Pm}}(\text{l})(298.15 \text{ K}))(T_{\text{fus}} - 298.15) & \quad (3) \quad 290 \end{aligned}$$

The terms $C_{\text{Pm}}(\text{s})(298.15 \text{ K})$ and $C_{\text{Pm}}(\text{l})(298.15 \text{ K})$ 291
 refer to the molar heat capacities of the solid and 292
 liquid state, respectively, of each ketone estimated at 293
 298.15 K by group additivity [9]. The results are re- 294

Table 9

A summary of the fusion, vaporization and sublimation enthalpies of ketones **1** and **2**

Compound	$C_{Pm}(l)$ (J mol ⁻¹)	$C_{Pm}(s)$ (J mol ⁻¹)	$\Delta_{fus}H_m(T_{fus})$ (kJ mol ⁻¹)	T_{fus} (K)	$\Delta_{fus}H_m(298.15\text{ K})$ (kJ mol ⁻¹)	$\Delta_{vap}H_m(298.15\text{ K})$ (kJ mol ⁻¹)	$\Delta_{sub}H_m(298.15\text{ K})$ (kJ mol ⁻¹)
Ketone 1	364.8	275.8	17.2 ± 0.5	366.6	12.8 ± 1.8	90.6 ± 2.0	103.3 ± 3.2
Ketone 2	364.8	275.8	27.8 ± 0.7	420	20.1 ± 2.9	92.0 ± 2.9	112.8 ± 4.1

ported in Table 9 and will be discussed below. Uncertainties represent two standard deviations of the mean. The uncertainty associated with this term was arbitrarily assigned as a fraction (0.167) of the total temperature adjustment.

3.3. Sublimation enthalpy

The sublimation enthalpies of ketones **1** and **2** are summarized in Table 9. The difference in the sublimation enthalpies of the two ketones is mainly due to the difference in fusion enthalpy.

3.4. Enthalpies of formation

The standard enthalpies of formation in the solid and ideal gas states were calculated using the data in Tables 9 and 10. In Table 10, experimental values of these quantities are shown together with values estimated by means of the Benson group additivity method. The values reported by the Benson method are for strainless molecules, i.e. ring-corrections were

ignored. The parameters needed for the estimation were taken from the literature [16].¹ Also included in Table 10 are estimations of molar enthalpies of formation of ketones **3–6**.

3.5. Quantum mechanical calculations

Ab initio quantum chemical calculations² were carried out for geometry optimized structures of isomers **1** and **2** at the RHF/3-21G(d)//RHF/3-21G(d) level [18].³ Numerical vibrational frequencies at 298.15 K and unscaled zero-point energies were also computed at same calculational level to obtain thermally corrected total energies. The individual energy contributions (including 0–298 K rotational and translational corrections) and the final total energy calculations are depicted in Table 11. For completeness, calculational results on the two remaining isomeric (uniquely “biphenyloid”) dibenzocyclooctadienones—ketones **5** and **6**—are also included in Table 11.

4. Discussion

It may be concluded, by examining the differences between experimental and estimated enthalpies of formation of ketones **1** and **2** listed in Table 10, that both these ketones are stabilized by roughly 30–40 kJ mol⁻¹ through the annelation of two benzenoid rings to the cyclooctane ring. The quantum chemical computations of Table 11 also confirm the stability




Fig. 2. A plot of $\Delta_{vap}H_m(298.15\text{ K})$ against $\Delta_{sl}^n H_m(T_m)$ for mixture 3.

¹ Group values were taken from reference [16]; a group value of -19.2 kJ mol^{-1} for $C-(H)_2(C_B)_2$ was derived from the Benson group values [16] and the gas phase molar enthalpy of formation of diphenylmethane ($-\Delta_f H^0(g)$: $165 \pm 1.5\text{ kJ mol}^{-1}$) [20].

² More precisely, we used the SPARTAN SGI Version 5.0.3 Open GL built under IRIX 6.2 by Wavefunction Inc., Irvine, CA, 1997 program for our quantum chemical calculations.

³ For a more thorough description of our calculational level, see [18].

Table 10

Standard molar enthalpies of sublimation and formation of ketones **1** and **2** in the gas phase at 298.15 K; comparisons with estimations by group additivity

Compound	$\Delta_f H^0(\text{s})$ (kJ mol ⁻¹)	$\Delta_{\text{sub}} H_m$ (kJ mol ⁻¹)	$\Delta_f H^0(\text{g})$ (kJ mol ⁻¹)	$\Delta_f H^0(\text{g})$ (kJ mol ⁻¹) Benson ^a
Ketone 1	-143.2 ± 4.3	103.3 ± 3.2	-39.9 ± 5.5	2.6
Ketone 2	-127.6 ± 3.7	112.8 ± 4.1	-14.8 ± 5.3	18.3
Ketone 3				4.5
Ketone 4				6.9
Ketone 5				2.0
Ketone 6				-1.5

^a Calculated from Benson group values [16] ignoring ring strain and using a group value of -19.2 kJ mol⁻¹ for C-(H)₂(CB)₂ [20].

order of ketones **1** and **2** obtained experimentally. The calculated difference ($\Delta\Delta_f H$) of 10 kJ mol⁻¹ is in relatively good agreement with the experimental difference of 24.1 ± 10.8 kJ mol⁻¹. This gives us confidence in our theoretical predictions for the other isomeric dibenzocyclooctadienones. With the exceptions of ketones **1** and **3**, the ab initio results shown in Table 11 are also in good relative agreement with the empirical estimations obtained by group additivity.

The additional stabilization observed here relative to those estimated by group methods, agree with previous results reported for dibenzoannelated cycloheptane derivatives [1,2]. For example, considering the seven membered ring homologue of ketone **2**, dibenzosuberone (2,3:6,7-dibenzocyclohepta-2,6-dien-1-one), a value of about 35 kJ mol⁻¹ has been reported for

the additional stabilization of this compound, due to the attachment of two benzenoid rings to the saturated carbocycle. A value exceeding 30 kJ mol⁻¹ was reported by Verevkin [17] for anthrone with its central six-membered ring.

Examination of the relative energies calculated by ab initio methods for the dibenzocyclooctadienones in Table 11 reveals that ketone **4** is stabilized anywhere from ~12 to 24 kJ mol⁻¹ compared to isomeric ketones **1** and **2**, even though the carbonyl groups in ketones **1** and **2** are conjugated to benzene rings. The strain energy associated with a sufficient flattening of the ring necessary for conjugation may be large enough that the ring remains non-planar at the cost of conjugation. Loss of conjugation of the carbonyl with the benzene ring can increase the ground state energy of the molecule because of the inductive

Table 11

RHF/3-21G(d)//RHF/3-21G(d) energies of isomeric dibenzocyclooctadienones; a comparison of ab initio results with those estimated by group additivity

Compound	E_{Total} (hartree) ^a	ZPE (kJ mol ⁻¹) ^b	Therm. corr. (kJ mol ⁻¹) ^c	E_{298} (hartree) ^d	$\Delta_f H_{\text{rel}}$ (kJ mol ⁻¹) ^e ab initio	$\Delta\Delta_f H_{\text{rel}}^{\text{rel}}$ (kJ mol ⁻¹) ^e Benson ^f
Ketone 1	-685.07083	726.4	757.7	-684.78223	31.2	4.1
Ketone 2	-685.06697	726.0	757.6	-684.77842	41.2	19.8
Ketone 3	-685.07324	727.6	758.3	-684.78441	25.5	6.0
Ketone 4	-685.07562	726.2	756.9	-684.78733	17.8	8.4
Ketone 5 ^g	-685.07676	724.3	753.9	-684.78961	11.8	3.5
Ketone 6	-685.08311	728.3	758.7	-684.79412	(0.0)	(0.0)

^a Total energies.

^b Zero-point energies.

^c Thermal corrections for translational, rotational, and vibrational enthalpy changes from 0 to 298 K.

^d Thermally corrected total energies, where $E_{298} = E_{\text{Total}} + \text{therm. corr.}$ and 1 Hartree = 2625.5 kJ mol⁻¹.

^e Calculated relative heats of formation, where (0.0) refers, by definition, to the lowest energy species.

^f Reference [16].

^g One imaginary frequency of 53.851 cm⁻¹ was ignored.


Table 12
Bond lengths (Å) measured for ketone **1**^a

Bond	Å
C(1)–C(6)	1.3967(16)
C(1)–C(7)	1.5125 (17)
C(2)–C(8)	1.5134(15)
C(4)–C(5)	1.3803(19)
C(7)–C(8)#1	1.5275(17)
C(8)–C(7)#1	1.5275(17)
C(1)–C(2)	1.4010(17)
C(2)–C(3)	1.3928(17)
C(3)–C(4)	1.3832(18)
C(5)–C(6)	1.3803(18)
C(8)–O(1)	1.0875

^a Symmetry transformations used to generate equivalent atoms: #1 – x, y, –z + 1/2.

effect of the non-conjugated phenyl ring on the electron deficient carbonyl carbon. Given similar losses in conjugation, this destabilizing effect would be expected to be greater in ketone **2** than in ketone **1**. The experimental data appears to bear this out. Ketone **1** is more stable than ketone **2**.

When this qualitative explanation is subjected to the other available experimental evidence, the data appears to be inconclusive. An examination of the crystal structure of ketone **1** reveals a dihedral angle of 28.3° between the carbonyl group and carbons C2–C3 of the benzene ring. To the extent that the solid state models the gas phase structure, conjugation should be attenuated by this angle. The bond length of the carbonyl group is not reliable because of disorder due to forced crystallographic symmetry (see Fig. 1, Table 12). An examination of the bond lengths between C(2)–C(8) (and crystallographically equivalent C(2)#1–C(8)#1 (see Table 12)) and C(1)–C(7) reveals identical distances between the four atoms. This distance between the benzene ring and the carbonyl group is an average of C(2)–C(8) and C(2)#1–C(8)#1. Nevertheless, the uncertainty associated with this bond length is very similar to the others in the crystal structure, suggesting both C(2)–C(8) and C(2)#1–C(8)#1 bond lengths to be very similar. An examination of crystal structures of 14 other *ortho* substituted conjugated aromatic ketones in the Cambridge Database with the following structure motif resulted in an average bond distance of (1.498 ± 0.03) Å (±2σ).



This compares to a value of (1.5134 ± 0.03) Å found in ketone **1**. Conjugation between the ring and the carbonyl group in the solid state of **1** on the basis of this argument would be expected to be attenuated. In contrast, the infrared frequency of 1681 cm⁻¹ (KBr) exhibited by ketone **1** in the solid state when compared to the frequency observed in acyclic 1,6-diphenylhexane-1,6-dione (ν_{max} 1681 cm⁻¹ (Nujol) [20]), appears normal for a ketone conjugated to a phenyl group. The infrared frequency for ketone **2** (1633 cm⁻¹ (CCl₄)), can be compared to the frequency observed in 4-phenylbenzophenone (1648 cm⁻¹ (Nujol) [20]). The crystal structure of ketone **2** is not available; thus any correlation of bond length with infrared frequency must await further experimental structural or gas phase studies.

Uncited reference

[19].

Acknowledgements

The authors express their thanks to Drs. L. Pârvolescu, A. Banciu, C. Costea and M. Pop for their valuable assistance in the synthesis of ketones **1** and **2**. JFL would like to thank Dow Chemical Co. for partial support of his thermochemical studies. JSC would like to thank NASA Glenn Research Center for financial support. DH would like to thank Fundação para a Ciência e a Tecnologia (Portugal) for a research fellowship (SFRH/BPD/5593/2001).

References

- [1] St. Perisanu, A. Banciu, M. Banciu, Thermochim. Acta 228 (1993) 393.
- [2] St. Perisanu, M. Banciu, Thermochim. Acta 320 (1998) 9.

- 436 [3] E. Cioranescu, A. Bucur, M. Elian, M. Banciu, M. Voicu, 454
437 C.D. Nenitzescu, *Tetrahedron Lett.* (1964) 3825. 455
- 438 [4] E. Cioranescu, A. Bucur, M. Banciu, C.D. Nenitzescu, *Rev.* 456
439 *Roum. Chimie* 10 (1965) 141. 457
- 440 [5] M. Banciu, C. Costea, D. Mihaiescu, *Rev. Roum. Chimie* 44 458
441 (1999) 867. 459
- 442 [6] M. Banciu, L. Părvulescu, A. Banciu, A. Simion, C. Costea, 460
443 C. Draghici, M. Pop, D. Mihaiescu, *J. Anal. Appl. Pyrolysis* 461
444 57 (2001) 261. 462
- 445 [7] R.H. Blessing, *Acta Cryst.* A51 (1995) 33–38. 463
- 446 [8] R.M. Stephenson, S. Malanowski, *Handbook of the Thermo-* 464
447 *dynamics of Organic Compounds*, Elsevier, New York, 1987 465
448 (vaporization enthalpies were calculated from the Antoine 466
449 constants). 467
- 450 [9] J.S. Chickos, D.G. Hesse, J.F. Liebman, *Struct. Chem.* 4 468
451 (1993) 261. 469
- 452 [10] J.S. Chickos, S. Hosseini, D.G. Hesse, J.F. Liebman, *Struct.* 470
453 *Chem.* 4 (1993) 271. 471
- [11] D.I. Marchidan, C. Telea, *Studii Cerc. Chimie* 3 (1971) 283. 454
- [12] I. Contineanu, C. Telea, *Rev. Chimie (Bucuresti)* (2001) 576; 455
I. Contineanu, C. Telea, *Romanian Patent No. 11* (1996). 456
- [13] The NBS tables of chemical thermodynamic properties, *J.* 457
Phys. Chem. Ref. Data (Suppl. 2) (1982) 11. 458
- [14] J.S. Chickos, D.G. Hesse, S. Hosseini, G. Nichols, P. Webb, 459
Thermochim. Acta 313 (1998) 101. 460
- [15] J.S. Chickos, *Thermochim. Acta* 313 (1998) 19. 461
- [16] S.W. Benson, *Thermochemical Kinetics*, 2nd ed., Wiley/
Interscience, New York, 1976. 462
- [17] S.P. Verevkin, *Thermochim. Acta* 310 (1998) 229. 463
- [18] W.J. Hehre, L. Radom, P.V.R. Schleyer, J.A. Pople, *Ab Initio* 464
Molecular Orbital Theory, Wiley, New York, 1986. 465
- [19] *Aldrich Library of FTIR Spectra*, 1st ed., Aldrich Chemical 466
Company. 467
- [20] (a) W.V. Steele, R.D. Chirico, N.K. Smith, *J. Chem. Ther-* 468
modyn. 27 (1995) 671; 469
(b) S.P. Verevkin, *J. Chem. Eng. Data* 44 (1999) 175. 470