



## The enthalpy of sublimation of cubane

A. Bashir-Hashemi<sup>a,1</sup>, James S. Chickos<sup>b,\*</sup>, William Hanshaw<sup>b</sup>, Hui Zhao<sup>b</sup>,  
Behzad S. Farivar<sup>c</sup>, Joel F. Liebman<sup>c</sup>

<sup>a</sup> Fluorochem, Inc., Azusa, CA 91702, USA

<sup>b</sup> Department of Chemistry and Biochemistry, University of Missouri at St. Louis, St. Louis, MO 63121, USA

<sup>c</sup> Department of Chemistry and Biochemistry, University of Maryland, Baltimore County, Baltimore, MD 21250, USA

Received 1 March 2004; received in revised form 6 May 2004; accepted 8 May 2004

This paper is dedicated to Philip E. Eaton and Thomas W. Cole, Jr. on the occasion of the 40th anniversary  
of the first successful synthesis of cubane and any of its derivatives

### Abstract

The sublimation enthalpy of cubane, a key reference material for force field and quantum mechanical computations, was measured by combining the vaporization enthalpy at  $T = 298.15$  K to the sum of the fusion enthalpy measured at  $T = 405$  K and a solid–solid phase transition that occurs at  $T = 394$  K. The fusion and solid–solid phase transitions were measured previously. A sublimation enthalpy value of  $(55.2 \pm 2.0)$  kJ mol<sup>-1</sup> at  $T = 298.15$  K was obtained. This value compares quite favorably the value obtained by comparing the sublimation enthalpy of similar substances as a function of their molar masses but is at odds with earlier measurements.

© 2004 Published by Elsevier B.V.

**Keywords:** Cubane; Sublimation; Enthalpy

### 1. Introduction

This is the 40th anniversary of the first successful synthesis of the polycyclic hydrocarbon cubane [1]: two other syntheses soon followed [2]. As befits the high symmetry and accompanying esthetics, large strain energy and thus high energy, and eight tertiary carbons all capable of possible functionalization, the chemistry of this seemingly simple 8-carbon hydrocarbon and its derivatives has blossomed as evidenced by numerous reviews in which it is featured prominently [3]. The eponymic (i.e., cubical) symmetry of cubane results in there being a single type of carbon environment and of bonded hydrogen, one type of C–C and C–H bond and associated bond lengths, and one unique C–C–C and C–C–H angle. Very few hydrocarbons have

such a minimal description<sup>2</sup>. Accordingly, paralleling the “organic” chemistry (and related bio- and high energy chemistry) interest in cubane and its derivatives, the physical chemists have been active—soon after the first synthesis of cubane itself there was a measurement of the enthalpies of combustion and of sublimation of this hydrocarbon [4] from which the gas phase enthalpy of formation of  $622.2 \pm 4.2$  kJ mol<sup>-1</sup> was derived. In turn, this quantity and the molecular high symmetry have meant that molecular mechanicians have been active: for example, cubane has been important in the development of the recent molecular mechanical model, MM4 [5a] and its predecessor, MM3 [5b] and quantum chemists have likewise been active with high

<sup>2</sup> The other known minimal hydrocarbons are methane, ethane, ethylene, acetylene, cyclopropane, neopentane, benzene, and dodecahedrane. Cyclohexane and cyclobutane, for example, do not qualify because there is a difference between equatorial and axial hydrogens; allene does not qualify because of two types of carbon, and cyclooctatetraene does not qualify because of two different types of carbon–carbon bonds. Ideally, diamond, graphite and polyethylene would qualify, however, there are end effects such as the finiteness of the sample.

\* Corresponding author. Tel.: +1 314 516 5377; fax: +1 314 516 5342.  
E-mail address: jsc@umsl.edu (J.S. Chickos).

<sup>1</sup> Present address: ERC Inc. at AFRL/PRS, 10 East Saturn Boulevard, Edwards AFB, CA 93524, USA.

level methodologies [6a,b]. Calculations at the G2(MP2) level give 606.7 or 625.9 kJ mol<sup>-1</sup> depending on whether the atomization or bond separation method is used [6a], while the G3(MP2) result is 610.9 kJ mol<sup>-1</sup>, 11.3 kJ mol<sup>-1</sup> lower than the reported experimental value [6b,c].

Recently using an analysis based on the sublimation enthalpies of other cyclic and polycyclic hydrocarbons, it has been suggested that the enthalpy of sublimation of cubane is seriously in error [7]. This raises considerable concern as to the enthalpy of formation of gaseous cubane and all related analyses. We recall that there is some controversy about the enthalpy of combustion, and thus formation of its

1,4-dicarbomethoxy derivative [8]. The current study reports a new experimental determination of the enthalpy of sublimation of cubane. Let us summarize our findings and analyses that follow: the just enunciated literature suggestion is verified and the derived concern is justified.

## 2. Experimental

All standards were purchased from the Aldrich Chemical Company and were used without any further purification. Each was analyzed by gas chromatography and found

Table 1  
Cubane mixture

CH <sub>2</sub> Cl <sub>2</sub>	t <sub>r</sub> (min)						
	348.7	353.9	358.9	364	369.1	374.2	
<b>(A) Mixture 1</b>							
Methane <sup>a</sup>	1.328	1.349	1.353	1.373	1.341	1.359	
Nonane	3.87	3.105	2.826	2.621	2.449	2.312	
Cubane	5.485	4.25	3.799	3.453	3.168	2.935	
Decane	6.964	5.081	4.437	3.927	3.524	3.199	
<i>exo</i> -THDCPD <sup>b</sup>	10.105	7.271	6.286	5.495	4.865	4.35	
Undecane	13.53	9.102	7.645	6.488	5.594	4.88	
<i>endo</i> -THDCPD <sup>b</sup>	12.247	8.658	7.404	6.488	5.645	4.993	
<b>(B) Mixture 2</b>							
	343.6	348.6	353.8	358.8	363.9	368.9	374
Methane <sup>a</sup>	1.27	1.283	1.29	1.342	1.344	1.357	1.375
Norbornene	1.957	1.879	1.809	1.76	1.71	1.68	1.665
Octane	2.481	2.302	2.154	2.04	1.947	1.875	1.83
Cubane	5.472	4.779	4.215	3.77	3.4	3.113	2.895
Adamantane	11.43	9.585	8.124	6.967	6.049	5.315	4.929
Undecane	13.833	11.199	9.207	7.662	6.54	5.59	4.929
Naphthalene	21.913	17.759	14.562	12.06	10.17	8.624	7.485
Dodecane	27.976	21.921	17.44	14.015	11.562	9.535	8.117
<b>(C) Mixture 3</b>							
	343.7	348.8	353.85	359	364	369.05	
Methane <sup>a</sup>	1.319	1.317	1.329	1.341	1.351	1.35	
<i>n</i> -Heptane	1.805	1.735	1.688	1.653	1.623	1.59	
Methylcyclohexane	1.961	1.872	1.809	1.76	1.718	1.675	
1-Octene	2.398	2.234	2.101	1.998	1.915	1.844	
Nonane	3.863	3.421	3.078	2.805	2.587	2.406	
Cubane	5.481	4.78	4.219	3.772	3.413	3.12	
<i>exo</i> -THDCPD <sup>b</sup>	10.491	8.994	7.494	6.423	5.594	5.012	
<i>endo</i> -THDCPD <sup>b</sup>	12.463	10.508	8.751	7.447	6.43	5.678	
	343.6	348.8	353.9	359.0	364	369.1	374.2
<b>(D) Mixture 4</b>							
	343.6	348.8	353.85	359.0	364	369.1	374.2
Methane <sup>a</sup>	1.295	1.301	1.295	1.304	1.315	1.326	1.325
<i>n</i> -Heptane	1.854	1.78	1.720	1.674	1.636	1.608	1.575
Methylcyclohexane	2.018	1.924	1.845	1.784	1.734	1.693	1.652
1-Octene	2.464	2.286	2.142	2.028	1.934	1.855	1.79
Nonane	3.952	3.495	3.128	2.84	2.609	2.421	2.263
Cubane	5.59	4.862	4.340	3.815	3.507	3.159	2.872
<i>exo</i> -THDCPD <sup>b</sup>	10.689	8.95	7.582	6.504	5.632	4.882	4.378
<i>endo</i> -THDCPD <sup>b</sup>	12.715	10.654	8.868	7.54	6.478	5.593	4.946

<sup>a</sup> On-retained reference.

<sup>b</sup> Tetrahydrodicyclopentadiene.

69 to be at least 99 mole percent pure. Cubane (+99 mol%)  
 70 was kindly supplied by Professor Phillip Eaton. Correla-  
 71 tion gas chromatography experiments were performed on  
 72 an HP 5890A Series II Gas Chromatograph equipped with  
 73 a split/splitless capillary injection port and a flame ioniza-  
 74 tion detector run at a split ratio of 100/1. Retention times  
 75 were recorded to three significant figures following the deci-  
 76 mal point on a HP 3989A Integrator. The instrument was  
 77 run isothermally using either a 15 or 30 m SPB-5 capillary  
 78 column. Helium was used as the carrier gas. At the temper-  
 79 atures of the experiments, the retention time of the solvent,  
 80  $\text{CH}_2\text{Cl}_2$ , decreased with increasing temperature suggesting  
 81 that it was retained by the column. Methane was bubbled  
 82 prior to each run and its retention time was found to in-  
 83 crease with temperature. A consequence of the increase in  
 84 viscosity of the carrier gas with temperature, this is the cri-  
 85 terion used to confirm that a substance is not retained on  
 86 the column and can be used to determine the dead volume  
 87 of the column. The gas chromatographic retention times of  
 88 cubane and the standards are summarized in Table 1. Ad-  
 89 justed retention times,  $t_a$ , were calculated by subtracting the  
 90 measured retention time of methane from the retention time  
 91 of each analyte as a function of temperature usually over a  
 92 30 K range. Column temperatures were controlled by the gas  
 93 chromatograph and were monitored independently by using  
 94 a Fluke 51 K/J thermometer. Temperature was maintained  
 95 constant by the gas chromatograph to  $\pm 0.1$  K.

### 96 3. Results

97 Cubane was analyzed using a series of different standards  
 98 in four separate mixtures. The literature values of the stan-  
 99 dards are reported in Table 2. Standards were chosen on the  
 100 basis of their relative retention times, our assessment of the  
 101 reliability of their vaporization enthalpies and their struc-  
 102 tural similarities to cubane. A plot of  $\ln(1/t_a)$  against  $1/T(\text{K})$   
 103 resulted in straight lines characterized by the parameters

Table 2

Literature values used as reference for cubane; molar enthalpies in  $\text{kJ mol}^{-1}$   $\Delta_{\text{vap}}H_m(298.15 \text{ K})$

	$\Delta_{\text{vap}}H_m(298.15 \text{ K})$	Reference
Norbornene	$35.1 \pm 0.2$	[9]
Methylcyclohexane	35.4	[10]
Heptane	$36.57 \pm 0.18$	[11]
1-Octene	$40.3 \pm 0.2$	[12]
Octane	$41.56 \pm 0.2$	[11]
Nonane	$46.55 \pm 0.46$	[11]
Adamantane	48.2	[13]
<i>exo</i> -Tetrahydrodicyclopentadiene	$49.1 \pm 2.3$	[14]
<i>endo</i> -Tetrahydrodicyclopentadiene	$50.2 \pm 2.3$	[14]
Decane	$51.42 \pm 0.26$	[11]
Naphthalene	$55.65 \pm 2.8$	[15]
Undecane	$56.58 \pm 0.56$	[11]
Dodecane	$61.52 \pm 0.61$	[11]




Fig. 1. The correlation between experimental and calculated vaporization enthalpies of the standards used in four separate correlations. The equation of the line calculated by a linear regression analysis is given by:  $\Delta_{\text{vap}}H_m(\text{kJ mol}^{-1}) (\text{lit}) = (1.004 \pm 0.034\Delta_{\text{vap}}H_m(\text{calcd}) - (0.15 \pm 0.98))$ .

104 listed in the second and third columns of Table 3. Equations  
 105 for the correlation of enthalpies of transfer from solution  
 106 to the vapor,  $\Delta_{\text{sln}}^v H_m(T_m)$ , against experimental vaporiza-  
 107 tion enthalpies are given at the bottom of Table 3 for each  
 108 correlation. A graphical summary of how well experimental  
 109 vaporization enthalpies were reproduced is given in Fig. 1.  
 110 The equation describing the correlation between experimen-  
 111 tal and calculated values of  $\Delta_{\text{vap}}H_m(298.15 \text{ K})$  is provided  
 112 in the caption of Fig. 1. The mean vaporization enthalpy  
 113 of cubane at  $T = 298.15 \text{ K}$  resulting from the four separate  
 114 correlations is  $(44.7 \pm 1.6) \text{ kJ mol}^{-1}$  (Table 4).

115 Solid phase transitions of cubane have been previously  
 116 measured by adiabatic calorimetry and DSC [17]. Two  
 117 phase transitions have been observed in the solid state of  
 118 cubane, a solid–solid transition at  $T_{\text{tr}} = (394.02 \pm 0.04) \text{ K}$   
 119 ( $\Delta_{\text{tr}}H_m(5.94 \pm 0.02) \text{ kJ mol}^{-1}$ ) measured by adiabatic  
 120 calorimetry and  $T_{\text{fus}}(\text{onset}) = (404.9 \pm 0.5) \text{ K}$  ( $\Delta_{\text{fus}}H_m(8.7$   
 121  $\pm 0.3) \text{ kJ mol}^{-1}$ ) measured by DSC [17]. Since both of  
 122 these transitions occur above  $T = 298.15 \text{ K}$ , both must be  
 123 taken into account in calculating the sublimation enthalpy of  
 124 cubane at  $T = 298.15$  using the following thermodynamic  
 125 equality:

$$\begin{aligned} \Delta_{\text{sub}}H_m(298.15 \text{ K}) \\ = \Delta_{\text{vap}}H_m(298.15 \text{ K}) + \Delta_{\text{cr}(2)}^1 H_m(298.15 \text{ K}) \end{aligned} \quad (1) \quad 127$$

129 where  $\Delta_{\text{vap}}H_m(298.15 \text{ K})$  represents the vaporization en-  
 130 thalpy at  $T = 298.15 \text{ K}$  and  $\Delta_{\text{cr}(2)}^1 H_m(298.15 \text{ K})$  represents  
 131 the sum of the measured phase transition and fusion enthalpy  
 132 adjusted to  $T = 298.15 \text{ K}$ .

133 The vaporization enthalpy of cubane at  $T = 298.15 \text{ K}$  is  
 134 obtained directly by correlation gas chromatography. Ad-

Table 3

	$\Delta_{\text{sln}}^{\text{v}} H_{\text{m}}(361 \text{ K})/R$	Intercept	$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K})$ (lit)	$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K})$ (calc)
(A) Mixture 1 <sup>a</sup>				
Cubane	4086.2	10.477		45.6
Nonane	4130.5	11.102	46.55	46.0
<i>exo</i> -THDCPD <sup>b</sup>	4533.4	11.028	49.1	49.7
<i>endo</i> -THDCPD <sup>b</sup>	4617.7	11.054	50.2	50.5
Decane	4713.1	11.996	51.42	51.4
Undecane	5232.7	12.733	56.58	56.2
(B) Mixture 2 <sup>c</sup>				
	$\Delta_{\text{sln}}^{\text{v}} H_{\text{m}}(359 \text{ K})/R$	Intercept	$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K})$ (lit)	$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K})$ (calc)
Norbornene	3772.8	11.351	35.1	36.4
Octane	4214.6	12.077	41.56	42.0
Cubane	4332.1	11.176		43.6
Adamantane	4539.4	10.909	48.2	46.2
Naphthalene	5163.4	12.008	55.65	54.3
Undecane	5358.1	13.074	56.58	56.8
Dodecane	5835.5	13.71	61.52	62.9
(C) Mixture 3 <sup>d</sup>				
	$\Delta_{\text{sln}}^{\text{v}} H_{\text{m}}(356 \text{ K})/R$	Intercept	$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K})$ (lit)	$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K})$ (calc)
Methylcyclohexane	3417.9	10.389	35.40	35.1
<i>n</i> -Heptane	3544.4	11.036	36.57	36.5
1-Octene	3951.2	11.421	40.3	41.0
Cubane	4292.3	11.065		44.7
Nonane	4411.4	11.904	46.55	46.1
<i>exo</i> -THDCPD <sup>b</sup>	4693.2	11.436	49.10	49.1
<i>endo</i> -THDCPD <sup>b</sup>	4793.4	11.535	50.20	50.2
(D) Mixture 4 <sup>e</sup>				
	$\Delta_{\text{sln}}^{\text{v}} H_{\text{m}}(356 \text{ K})/R$	Intercept	$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K})$ (lit)	$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K})$ (calc)
Methylcyclohexane	3346.6	10.063	35.40	35.7
<i>n</i> -Heptane	3385.7	10.434	36.57	36.1
1-Octene	3898.0	11.19	40.3	41.0
Cubane	4256.3	10.927		44.4
Nonane	4386.0	11.789	46.55	45.7
<i>exo</i> -THDCPD <sup>b</sup>	4763.0	11.622	49.10	49.3
<i>endo</i> -THDCPD <sup>b</sup>	4870.3	11.736	50.20	50.3

<sup>a</sup>  $\Delta H_{\text{vap}}(298.15 \text{ K}) \text{ kJ mol}^{-1} = (1.117 \pm 0.085) \Delta_{\text{sln}}^{\text{v}} H_{\text{m}}(361 \text{ K}) + (7.63 \pm 0.56)$ ;  $r^2 = 0.9829$ .

<sup>b</sup> Tetrahydrodicyclopentadiene.

<sup>c</sup>  $\Delta_{\text{I}}^{\text{g}} H_{\text{m}}(298.15 \text{ K}) \text{ kJ mol}^{-1} = (1.55 \pm 0.11) \Delta_{\text{sln}}^{\text{v}} H_{\text{m}}(359 \text{ K}) - (12.24 \pm 1.55)$ ;  $r^2 = 0.9809$ .

<sup>d</sup>  $\Delta_{\text{I}}^{\text{g}} H_{\text{m}}(298.15 \text{ K}) \text{ kJ mol}^{-1} = (1.321 \pm 0.041) \Delta_{\text{sln}}^{\text{v}} H_{\text{m}}(359 \text{ K}) - (2.39 \pm 0.45)$ ;  $r^2 = 0.9961$ .

<sup>e</sup>  $\Delta_{\text{I}}^{\text{g}} H_{\text{m}}(298.15 \text{ K}) \text{ kJ mol}^{-1} = (1.157 \pm 0.052) \Delta_{\text{sln}}^{\text{v}} H_{\text{m}}(359 \text{ K}) + (3.52 \pm 0.64)$ ;  $r^2 = 0.9921$ .

Table 4

Summary of vaporization enthalpies; by correlation and from the literature

	$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K})$ (lit)	Mix 1	Mix 2	Mix 3	Mix 4	$\Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K})$ mean
Norbornene	35.1		36.4			36.4
Methylcyclohexane	35.4			35.1	35.7	35.4
Heptane	36.57			36.5	36.1	36.3
1-Octene	40.3			41.0	41.0	41.0
Octane	41.56		42.0			42.0
Cubane		45.6	43.6	44.7	44.4	44.6 ± 0.8
Nonane	46.55	46.0		46.1	45.7	45.9
Adamantane	48.2		46.2			46.2
<i>exo</i> -THDCPD <sup>a</sup>	49.1	49.7		49.1	49.3	49.2
<i>endo</i> -THDCPD <sup>a</sup>	50.2	50.5		50.2	50.3	50.3
Decane	51.42	51.4				51.4
Naphthalene	55.65		54.3			54.3
Undecane	56.58	56.2	56.8			56.5
Dodecane	61.52		62.9			62.9

<sup>a</sup> Tetrahydrodicyclopentadiene.

Table 5  
Phase change enthalpies of cubane; enthalpies in  $\text{kJ mol}^{-1}$

$\Delta_{\text{cr}(2)}^{\text{cr}(1)} H_{\text{m}}(394 \text{ K})$ (lit) <sup>a</sup>	$5.94 \pm 0.02$
$\Delta_{\text{cr}(1)}^{\text{cr}(1)} H_{\text{m}}(T_{\text{fus}})$ (lit) <sup>a</sup>	$8.7 \pm 0.3$
$T_{\text{fus}}$ (K)	404.9
$\Delta_{\text{cr}}^1 C_{\text{p,m}} \Delta T$ $\text{kJ mol}^{-1}$ <sup>b</sup>	$-4.02 \pm 1.2$
$\Delta_{\text{cr}(2)}^1 H_{\text{m}}(298 \text{ K})$	$10.6 \pm 1.2$
$\Delta_{\text{vap}} H_{\text{m}}(298 \text{ K})$ <sup>c</sup>	$44.6 \pm 1.6$
$\Delta_{\text{sub}} H_{\text{m}}(298 \text{ K})$	$55.2 \pm 2.0$

<sup>a</sup> [17].

<sup>b</sup> The experimental heat capacity of the crystal at  $T = 298.15 \text{ K}$  was obtained graphically [17]:  $C_{\text{p}}(\text{cr}) = 125 \text{ J mol}^{-1} \text{ K}^{-1}$ ; the heat capacity of the liquid phase of cubane was estimated [16]:  $C_{\text{p}}(\text{Q}(l)) = 179.2 \text{ J mol}^{-1} \text{ K}^{-1}$ ; the total phase change enthalpy,  $\Delta_{\text{cr}(2)}^1 H_{\text{m}}(T_{\text{fus}})$  was adjusted to  $T = 298.15 \text{ K}$  using Eq. (4) [18]; the uncertainty ( $\pm 2\sigma$ ) assumed to be 0.3 of the magnitude of the temperature adjustment, see [18] for further details.

<sup>c</sup> Uncertainty in vaporization enthalpy represents  $\pm 2\sigma$  of the mean.

justment of the total phase change enthalpy from  $T = T_{\text{fus}}$  to  $T = 298.15 \text{ K}$  is necessary because of the difference in heat capacity of the crystalline and liquid phases. A protocol for doing this is described below (Eqs. (4) and (5)); the method has recently been tested [18]. Inclusion of the temperature adjustment, which is small, results in a sublimation enthalpy for cubane at  $T = 298.15 \text{ K}$  of  $(55.2 \pm 2.0) \text{ kJ mol}^{-1}$ . These results are summarized in Table 5. This value is considerably smaller than the value of  $(80.3 \pm 1.7) \text{ kJ mol}^{-1}$  measured by Knudsen effusion reported previously [4].

#### 4. Discussion

The sublimation enthalpy of cubane has previously been calculated by the atom–atom potential method. The value calculated,  $62.8 \text{ kJ mol}^{-1}$  [19], is in reasonably good agreement with the value of  $(55.2 \pm 2.0) \text{ kJ mol}^{-1}$  obtained in this study. V.V. Diky et al. in their article questioning the sublimation enthalpy of cubane demonstrate that the sublimation enthalpies of a variety of saturated cyclic and polycyclic hydrocarbons correlate on a qualitative basis, with their molar mass. A graph similar to theirs is reproduced in Fig. 2 using the data in Table 6 obtained from recent compendia [20–22] and includes the sublimation enthalpy of cubane determined previously (●) and by this work (■). Literature vaporization and sublimation enthalpies, columns 2 and 3 of Table 6, were adjusted to  $T = 298.15 \text{ K}$  when necessary, using Eqs. (2) and (3), respectively. The sublimation enthalpy of the rigid solid,  $[\Delta_{\text{sub}} H_{\text{m}}(298.15 \text{ K})]$  column 10, Table 6, was calculated by combining the sublimation enthalpy with all solid–solid phase transitions occurring between  $T = 0 \text{ K}$ , and the temperature(s) at which the sublimation enthalpy was measured, columns 3 and 4. For compounds that are liquids at  $T = 298.15 \text{ K}$ , the vaporization enthalpy was adjusted to  $T = 298.15 \text{ K}$  using Eq. (2) when necessary and the fusion enthalpy was adjusted to  $T = 298.15 \text{ K}$  using Eq. (4) [16,18]. Their sum (columns 2 and 9), combined with any solid–solid phase transitions observed (column 4), are also




Fig. 2. Relation of enthalpies of sublimation of the rigid solid ( $[\Delta_{\text{sub}} H_{\text{m}}]$ ) at  $T = 298.15 \text{ K}$  to their molar mass,  $M$  [7]. Cage hydrocarbons ( $\square$ ); bicyclic compounds ( $\Delta$ ); cyclic compounds ( $\circ$ ). Cubane: ( $\blacksquare$ ) this work. Cubane: ( $\bullet$ ) lit. [4]. The solid line was obtained by a linear regression analysis.

listed in column 10. Sublimation enthalpies of the rigid solid were calculated in a slightly different manner by Diky et al. [7]; however, the results compare favorably. All heat capacities, in column 6 were estimated unless noted otherwise.

$$\begin{aligned} \Delta_{\text{vap}} H_{\text{m}}(298.15 \text{ K}) (\text{kJ mol}^{-1}) & \\ &= \Delta_{\text{vap}} H_{\text{m}}(T_{\text{m}}) + \frac{(10.58 + 0.26 \times C_{\text{p,l}})(T_{\text{m}} - 298.15)}{1000} \end{aligned} \quad (2)$$

$$\begin{aligned} \Delta_{\text{sub}} H_{\text{m}}(298.15 \text{ K}) (\text{kJ mol}^{-1}) & \\ &= \Delta_{\text{sub}} H_{\text{m}}(T_{\text{m}}) + \frac{(0.75 + 0.15 C_{\text{p,cr}})(T_{\text{m}} - 298.15)}{1000} \end{aligned} \quad (3)$$

$$\begin{aligned} \Delta_{\text{fus}} H_{\text{m}}(298.15 \text{ K}) / \text{kJ mol}^{-1} & \\ &= \Delta_{\text{fus}} H_{\text{m}}(T_{\text{fus}}) + \Delta_{\text{cr}}^1 C_{\text{p,m}} \Delta T, \end{aligned} \quad (4)$$

where

$$\begin{aligned} \Delta_{\text{cr}}^1 C_{\text{p,m}} \Delta T (\text{kJ mol}^{-1}) & \\ &= [0.15 C_{\text{p}}(\text{cr}) - 0.26 C_{\text{p}}(l) - 9.83] \left( \frac{T_{\text{fus}} - 298.15}{1000} \right). \end{aligned} \quad (5)$$

A treatment of the data in the graph by a linear regression analysis, excluding cubane from the analysis, results in the following relationship between sublimation enthalpy and molar mass ( $M$ ):

$$\begin{aligned} \Delta_{\text{sub}} H_{\text{m}}(298.15 \text{ K}) (\text{kJ mol}^{-1}) & \\ &= (0.425 \pm 0.023)M + (6.64 \pm 4.6); \quad r^2 = 0.9535 \end{aligned} \quad (6)$$

Table 6  
Enthalpies of sublimation of saturated cyclic hydrocarbons and auxiliary data<sup>a</sup>

	$\Delta_{\text{vap}}H_{\text{m}}$ (298 K)	$\Delta_{\text{sub}}H_{\text{m}}$ (298 K)	$\Delta_{\text{tr}}H_{\text{m}}$ ( $T_{\text{tr}}$ )	$T_{\text{tr}}$ (K)	Cp(l)/ Cp(cr)	$\Delta_{\text{fus}}H_{\text{m}}$ ( $T_{\text{fus}}$ )	$T_{\text{fus}}$ (K)	$\Delta_{\text{fus}}H_{\text{m}}$ (298 K)	$[\Delta_{\text{sub}}H_{\text{m}}]^{\text{b}}$ (298 K)	$M$ (g/mol)	$\Delta_{\text{sub}}H_{\text{m}}$ (298 K) Calc.
Cyclic compounds											
Cyclobutane	24.1		5.71	145.7	104/98.4	1.09	182.4	5.4	35.3	56.11	30.5
Cyclopentane	28.5		4.9	122	130/123						
			0.34	138		0.6	179.7	5.84	39.6	70.1	36.4
Cyclohexane	33.04		6.74	186.1	155/148	2.68	279.8	3.62	43.4	84.16	42.4
Cycloheptane	38.53		4.97	134.8							
			0.29	198.2							
			0.45	212.4	181/172	1.89	265.1	3.8	48.0	98.19	48.4
Cyclooctane	43.35		6.31	166.5	207/197						
			0.48	183.8		2.41	287.9	3.1	53.25	112.21	54.3
Cyclododecane		76.1	0.6	199					76.7	168.32	78.2
Cyclotetradecane		96.8							96.8	192.3	88.4
Bicyclic and tricyclic compounds											
Bicyclooctane		47.7	4.6	164.3			52.3	110.2	53.5		
<i>cis</i> -PHI <sup>c</sup>	46.12		8.26	182.3							
			0.39	184.5	223/196	1.4	236.5	3.3	58.1	124.2	59.4
<i>trans</i> -PHI <sup>c</sup>	44.76				223/196	10.9	213.9	14.1	58.9	124.2	59.4
<i>m</i> -Decalin	50.1		2.14	216.1	248/220	9.49	230.2	14.6	66.8	138.3	65.4
<i>trans</i> -Decalin	48.5				248/220	14.4	242.8	18.6	67.1	138.3	65.4
<i>endo</i> -THDCPD <sup>d</sup>		51	10.7	213.8					61.7	136.2	64.5
<i>exo</i> -THDCPD <sup>d</sup>	49.1		3.18	162.1	238/194	1.2	183.2	6.35	58.6	136.2	64.5
Cage compounds											
Cubane									55.2	104.2	50.9
Adamantane		59.3 <sup>e</sup>	3.38	208.7					62.6	136.2	64.5
PCU <sup>f</sup>		55.85	4.86						60.7	146.2	68.8
HCTD <sup>g</sup>		79.29							79.3	184.3	84.9
Diamantane		96.77							96.77	188.3	86.7
Pagodane <sup>h</sup>		116.8			/263				116.8	260.4	117.3

<sup>a</sup> Enthalpies in  $\text{kJ mol}^{-1}$ ; estimated heat capacities of the liquid/crystal (Cp(l)/Cp(cr) in  $\text{J mol}^{-1} \text{K}^{-1}$  [16]; phase change enthalpies were obtained from references [20–22] unless referenced otherwise.

<sup>b</sup>  $[\Delta_{\text{sub}}H_{\text{m}}](298) = \Delta_{\text{vap}}H_{\text{m}}(298) + \Delta_{\text{fus}}H_{\text{m}}(298) + \Sigma\Delta_{\text{tr}}H_{\text{m}}(T_{\text{tr}})$  or  $\Delta_{\text{sub}}H_{\text{m}}(298) + \Sigma\Delta_{\text{tr}}H_{\text{m}}(T_{\text{tr}})$ ; sublimation of the rigid crystal.

<sup>c</sup> Perhydroindane [7].

<sup>d</sup> Tetrahydrodicyclopentadiene [14].

<sup>e</sup> Average of six values reported at  $T = 298.16 \text{ K}$  [20].

<sup>f</sup> Pentacyclo[5.4.0.0<sup>2,6</sup>.0<sup>3,10</sup>.0<sup>5,9</sup>]undecane [26].

<sup>g</sup> Heptacyclo[6.6.0<sup>2,6</sup>.0<sup>3,13</sup>.0<sup>4,11</sup>.0<sup>5,9</sup>.0<sup>10,14</sup>]tetradecane [28].

<sup>h</sup> Adjusted to  $T = 298.15 \text{ K}$  using Eq. (3) using an estimated Cp(cr) of  $263 \text{ J mol}^{-1} \text{K}^{-1}$  [29].

198 The sublimation enthalpy of cubane calculates to  $(50.9$   
199  $\pm 4.6) \text{ kJ mol}^{-1}$  according to this equation. The results  
200 calculated using this equation are included as the last  
201 column in Table 6. The experimental sublimation en-  
202 thalpy of cubane is within the uncertainty of the corre-  
203 lation.

204 Using the group contribution method reported by Diky  
205 et al. [7] for estimating the sublimation enthalpy of the rigid  
206 crystalline state, a group value of  $(5.52 \pm 0.41) \text{ kJ mol}^{-1}$   
207 is derived for a cyclic tertiary  $\text{sp}^3$  carbon from bicyclic and  
208 polycyclic model compounds containing five and six mem-  
209 bered rings. This computes to a sublimation enthalpy of  
210  $(44.2 \pm 3.3) \text{ kJ mol}^{-1}$  estimated for cubane. The experimen-  
211 tal value determined is about  $10 \text{ kJ mol}^{-1}$  larger. Using the  
212 values reported in Table 6 to generate group values for cyclic  
213 tertiary and secondary  $\text{sp}^3$  hybridized carbon atoms result  
214 in near identical group values, 5.59 and  $6.87 \text{ kJ mol}^{-1}$ , re-  
215 spectively.

The sublimation enthalpy of cubane can also be estimated  
216 by combining an estimated vaporization enthalpy with the  
217 experimental fusion enthalpy. Using the following equation  
218 for predicting the vaporization enthalpy of a hydrocarbon  
219 [23]:  
220

$$\Delta_{\text{vap}}H_{\text{m}}(298.15 \text{ K}) (\text{kJ mol}^{-1}) = 4.69(n_{\text{C}} - n_{\text{Q}}) + 1.3n_{\text{Q}} + 3.0 \quad (7)$$

224 where  $n_{\text{C}}$  equals the number of carbon atoms and  $n_{\text{Q}}$  refers  
225 to the number of quaternary carbons, a vaporization en-  
226 thalpy of  $40.5 \text{ kJ mol}^{-1}$  results. Addition of the temperature  
227 adjusted fusion enthalpy of  $10.6 \text{ kJ mol}^{-1}$  result in an es-  
228 timated sublimation enthalpy of  $51.1 \text{ kJ mol}^{-1}$ , a value in  
229 good agreement with the experimental determination of  
230  $(55.2 \pm 2.0) \text{ kJ mol}^{-1}$ .

231 As noted by Diky et al. [7], the sublimation enthalpy  
232 of cyclotetradecane, also measured by Knudsen effusion by

233 the same laboratory reporting the cubane results [24], is  
 234 similarly much larger than results reported by others for the  
 235 same compound [25–27].

236 The enthalpy of formation of crystalline cubane [4], is  
 237 also subject to question. The value of  $541.8 \text{ kJ mol}^{-1}$  de-  
 238 rived from combustion measurements ( $\Delta_c E^\circ = -4828.3 \text{ kJ}$   
 239  $\text{mol}^{-1}$ ), was corrected for an unspecified amount of ill-  
 240 defined carbon adhering to the walls of the bomb. Unravel-  
 241 ing of this portion of the problem will also require additional  
 242 experimental measurements.

## 243 Acknowledgements

244 Financial support from the Research Board of the Uni-  
 245 versity of Missouri is gratefully acknowledged. We would  
 246 also like to thank Professors Phillip E. Eaton for a sample of  
 247 cubane Donald Rogers (Emeritus) for communicating the re-  
 248 sults of his computations and Vladimir Diky and his cowork-  
 249 ers for communicating their results prior to publication.

## 250 References

- 251 [1] P.E. Eaton, T.W. Cole, *J. Am. Chem. Soc.* 86 (1964) 962.  
 252 [2] (a) For almost contemporaneous alternative approaches to the syn-  
 253 thesis of cubane see: J.C. Barborak, L. Watts, R.J. Pettit, *J. Am.*  
 254 *Chem. Soc.* 88 (1966) 1328;  
 255 (b) C.G. Chin, W.H. Cuts, S. Masamune, *J. Chem. Soc. Chem. Com-*  
 256 *mun.* (1966) 880.  
 257 [3] (a) For some recent reviews see: G.W. Griffin, A.P. Marchand, *Chem.*  
 258 *Rev.* 89 (1989) 997;  
 259 (b) P.E. Eaton, *Angew. Chem. Int. Ed. Engl.* 31 (1992) 1421;  
 260 (c) H. Higuchi, I. Ueda, in: E.J. Osawa, O. Yonemitsu (Eds.), *Car-*  
 261 *bocyclic Cage Compounds; Chemistry and Applications*, vol. 217,  
 262 VCH, New York, NY, 1992;  
 263 (d) A. Bashir-Hashemi, S. Iyer, J. Alster, N. Slagg, *Chem. Ind.* 14  
 264 (1995) 551;  
 265 (e) J. Tsanakatsidis, in: B. Halton (Ed.), *Advances in Strain in Or-*  
 266 *ganic Chemistry*, vol. 6, JAI Press Inc., Greenwich, CT, 1997, p. 67;  
 267 (f) A. Bashir-Hashemi, in: K. Laali (Ed.), *Advances in Strained and*  
 268 *Interesting Organic Molecules*, JAI Press Inc., Greenwich, CT, 1999,  
 269 p. 1;  
 270 (g) A. Bashir-Hashemi, H. Higuchi, in: Z. Rappoport, J.F. Liebman  
 271 (Eds.) *The Chemistry of Cyclobutanes*, Wiley, Chichester (volume  
 272 nearing completion);  
 273 (h) E. Quintilla, J.Z. Davalos, J.L.M. Abboud, I. Alkorta, in: Z.  
 274 Rappoport, J.F. Liebman, (Eds.), *The Chemistry of Cyclobutanes*,  
 275 Wiley, Chichester (volume nearing completion).  
 276 [4] B.D. Kybett, S. Carroll, P. Natalis, D.W. Bonnell, J.L. Margrave,  
 277 J.L. Franklin, *J. Am. Chem. Soc.* 88 (1966) 626.  
 278 [5] (a) K.-H. Chen, N.L. Allinger, *J. Mol. Struct. (THEOCHEM)* 581  
 279 (2002) 215;  
 280 (b) N.L. Allinger, Y.H. Yuh, J.H. Lu, *J. Am. Chem. Soc.* 111 (1989)  
 281 8551.
- [6] (a) O. Castaño, R. Notario, J.-L.M. Abboud, R. Gomperts, L.-M. 282  
 Frutos, *J. Org. Chem.* 64 (1999) 9015; 283  
 (b) D.W. Rogers, *J. Mol. Struct.* 556 (2000) 207; 284  
 (c) The value reported in reference 6b is in error. The G3(MP2) result 285  
 is  $610.9 \text{ kJ mol}^{-1}$ ; personal communication with Professor Donald 286  
 W. Rogers. 287
- [7] V.V. Diky, M. Frenkel, L.S. Karpushenkava, *Thermochim. Acta* 408 288  
 (2003) 115. 289
- [8] (a) D.R. Kirklin, K.L. Churney, E.S. Domalski, *J. Chem. Thermodyn.* 290  
 21 (1989) 1105; 291  
 (b) V.V. Avdonin, E.I. Kipichev, Y.I. Rubtsov, L.B. Romanova, M.E. 292  
 Ivanova, L.T. Eremenko, *Russ. Chem. Bull.* 45 (1996) 2342. 293
- [9] W.V. Steele, R.D. Chirico, S.E. Knipmeyer, A. Nguyen, N.K. Smith, 294  
*J. Chem. Eng. Data* 41 (1996) 1285. 295
- [10] P.T. Eubank, L.E. Cedlel, J.C. Holste, K.R. Hall, *J. Chem. Eng. Data* 296  
 29 (1984) 389. 297
- [11] K. Ruzicka, V.J. Majer, *J. Phys. Chem. Ref. Data* 23 (1994) 1. 298
- [12] M. Mansson, P. Sellers, G. Stridh, S. Sunner, *J. Chem. Thermodyn.* 299  
 9 (1977) 91. 300
- [13] A. van Roon, J.R. Parsons, H.A.J. Govers, *J. Chromatogr. A* 955 301  
 (2002) 105. 302
- [14] J.S. Chickos, D.M. Hillesheim, G. Nichols, M.J. Zehe, *J. Chem.* 303  
*Thermodyn.* 34 (2002) 1647. 304
- [15] R. Sabbah, A. Xu-wu, J.S. Chickos, M.L. Planas Leitao, M.V. Roux, 305  
 L.A. Torres, *Thermochim. Acta* 331 (1999) 93. 306
- [16] J.S. Chickos, S. Hosseini, D.G. Hesse, J.F. Liebman, *Struct. Chem.* 307  
 4 (1993) 271; 308  
 J.S. Chickos, S. Hosseini, D.G. Hesse, J.F. Liebman, *Struct. Chem.* 309  
 4 (1993) 261. 310
- [17] M.A. White, R.E. Wasylshen, P.E. Eaton, Y. Xiong, K. Pramod, N. 311  
 Nodari, *J. Phys. Chem.* 96 (1992) 421. 312
- [18] J.S. Chickos, *Thermochim. Acta* 313 (1998) 19. 313
- [19] C.A. Fyfe, D. Harold-Smith, *J. Chem. Soc., Faraday Trans. II* 71 314  
 (1975) 4680. 315
- [20] J.S. Chickos, W.E. Acree Jr., *J. Phys. Chem. Ref. Data* 31 (2002) 316  
 537. 317
- [21] J.S. Chickos, W.E. Acree Jr., J.F. Liebman, *J. Phys. Chem. Ref.* 318  
*Data* 28 (1999) 1535. 319
- [22] J.S. Chickos, W.E. Acree Jr., *J. Phys. Chem. Ref. Data* 32 (2003) 320  
 515. 321
- [23] (a) J.S. Chickos, W.E. Acree Jr., J.F. Liebman, in: K.K. Irikura, D.J. 322  
 Frurip (Eds.), *Computational Thermochemistry*, ACS Symposium 323  
 Series 677, ACS, Washington DC, 1998 (Chapter 4); 324  
 (b) J.S. Chickos, A.S. Hyman, L.A. Ladon, J.F. Liebman, *J. Org.* 325  
*Chem.* 46 (1981) 4294. 326
- [24] M.A. Frisch, R.G. Bautista, J.L. Margrave, C.G. Parsons, J.H. Wotiz, 327  
*J. Am. Chem. Soc.* 86 (1964) 335. 328
- [25] J.S. Chickos, D.G. Hesse, S.Y. Panshin, D.W. Rogers, M. Saunders, 329  
 P.M. Uffer, J.F. Liebman, *J. Org. Chem.* 57 (1992) 1897. 330
- [26] G.J. Kabo, A.A. Kozyo, V.V. Diky, V.V. Simirsky, L.S. Ivashkevich, 331  
 A.P. Krasulin, V.M. Sevruk, M. Frenkel, A.P. Marchand, *J. Chem.* 332  
*Thermodyn.* 27 (1995) 707. 333
- [27] E. Eliel, J. Engelsman, *J. Chem. Ed.* 73 (1996) 903. 334
- [28] G.J. Kabo, A.A. Kozyo, A.P. Marchand, V.V. Diky, V.V. Simirsky, 335  
 L.S. Ivashkevich, A.P. Krasulin, V.M. Sevruk, M. Frenkel, *J. Chem.* 336  
*Thermodyn.* 26 (1994) 129. 337
- [29] H.-D. Beckhaus, C. Rtichardt, D.R. Lagerwall, L.A. Paquette, F. 338  
 Wahl, H. Prinzbach, *J. Am. Chem. Soc.* 116 (1994) 11775; 339  
 H.-D. Beckhaus, C. Rtichardt, D.R. Lagerwall, L.A. Paquette, F. 340  
 Wahl, H. Prinzbach, *J. Am. Chem. Soc.* 117 (1995) 8885. 341